1

(b)

(c)

Alpha, beta and gamma are types of nuclear radiation.

(a) Draw **one** line from each type of radiation to what the radiation consists of.

## What radiation consists of Type of radiation Electron from the nucleus Alpha Two protons and two neutrons Beta Electromagnetic radiation Gamma Neutron from the nucleus (3) A teacher demonstrates the penetration of alpha, beta and gamma radiation through different materials. The demonstration is shown in the figure below. Paper 5 mm of 5 cm of aluminium lead Complete the figure above by writing the name of the correct radiation in each box. (2) Give **two** safety precautions the teacher should have taken in the demonstration.

www.tutorzone.co.uk The table below shows how the count rate from a radioactive source changes with time. (d)

Time in seconds	0	40	80	120	160
Count rate in counts / second	400	283	200	141	100

	Use the table to calculate the count rate after 200 seconds.	
		(2)
(e)	The half-life of the radioactive source used was very short.	
	Give <b>one</b> reason why this radioactive source would be much less hazardous after 800 seconds.	
		(1)
	(Total 10 ma	arks)

Atoms contain three types of particle. 2

> (a) Draw a ring around the correct answer to complete the sentence.

The particles in the nucleus of the atom are

electrons and neutrons. electrons and protons. neutrons and protons.

(1)

Complete the table to show the relative charges of the atomic particles. (b)

Particle	Relative charge
Electron	-1
Neutron	
Proton	

www.tutorzone.co.uk

(c)	(i)	A neutral atom has no overall charge.	www.tutorzone.co.
		Explain this in terms of its particles.	
			(2)
	(ii)	Complete the sentence.	
		An atom that loses an electron is called an	
		and has an overall charge.	(2)

www.tutorzone.co.uk
In this question you will be assessed on using good English, organising information clearly

Some substances are radioactive. They may emit alpha or beta particles.  Describe the characteristics of alpha particles and beta particles in terms of their:  structure  penetration through air and other materials  deflection in an electric field.
<ul> <li>structure</li> <li>penetration through air and other materials</li> <li>deflection in an electric field.</li> </ul>
penetration through air and other materials     deflection in an electric field.
(6) (Total 13 marks)

(d)

$\mathbf{a}$	
J	

(b)

Nuclear fission and nuclear fusion are two processes that release energy.

<ul><li>(a) (i) Use the correct answer from the box to complete each sent</li></ul>	(a)	each sentence
---	-----	---------------

	Geiger counter	nuclear reactor	star	
	Nuclear fission takes place	e within a		
	Nuclear fusion takes place	e within a		
(ii)	State <b>one</b> way in which the nuclear fission.	e process of nuclear fusior	n differs from the	
				(1)
The	following nuclear equation r	represents the fission of ur	anium-235 (U-23	35).
	${}^{1}_{0}$ n + ${}^{235}_{92}$ U $\longrightarrow$ ${}^{236}_{92}$ U	> <sup>141</sup> <sub>56</sub> Ba + <sup>92</sup> <sub>36</sub> Kr	+ 3 <sup>1</sup> <sub>0</sub> n + ene	ergy
Chei	mical symbols:			
	Ba - barium			
	Kr - krypton			
(i)	Use the information in the	equation to describe the p	rocess of nuclea	r fission.
				•••

(4)

(ii) An isotope of barium is Ba-139. Ba-139 decays by beta decay to lanthanum-139 (La-139).


Complete the nuclear equation that represents the decay of Ba-139 to La-139.

(a) The names of three types of radiation are given in **List A**. Some properties of these three types of radiation are given in **List B**.

Draw **one** line from each type of radiation in **List A** to its correct property in **List B**.

List A Type of radiation	List B Property of radiation
	will pass through paper but is stopped by thin metal
alpha	
	has the shortest range in air
beta	
	will not harm human cells
gamma	
	is very weakly ionising

The radioactive isotope iodine-123 can be used by a doctor to examine the thyroid gland of a patient. The iodine, taken as a tablet in absorbed level of the control of the (b) a patient. The iodine, taken as a tablet, is absorbed by the thyroid gland. The gamma radiation emitted as the iodine atoms decay is detected outside the body.



The doctor uses an isotope emitting gamma radiation to examine the thyroid gland rather than an isotope emitting alpha or beta radiation.

Which one of the following gives a reason why gamma radiation is used?

Tick (✓) **one** box.

(c)

After 13 hours		0	f the iodine	1 e-123 atoms the thyroid absorbed	
	all	half	most		
Use a word from	the box to con	nplete the se	entence.		
lodine-123 has a	half-life of 13 l	nours.			
					(1)
Gamma radiation	ı has a long raı	nge in air.			
Gamma radiation	is not deflecte	ed by a magr	net.		
Gamma radiation	ı will pass throi	ugh the body	<i>1</i> .		

Page 7 of 99

(d) lodine-123 and iodine-131 are two of the isotopes of iodine.

Draw a ring around the correct answer to complete the sentence.

The nucleus of an iodine-123 atom has the same number of

electrons
neutrons as the
protons

nucleus of an iodine-131 atom.

(1) (Total 6 marks)

In 2011 an earthquake caused severe damage to a nuclear power station in Japan.

The damage led to the release of large amounts of radioactive iodine-131  $\binom{131}{53}$ I) into the atmosphere.

(a) The table gives some information about an atom of iodine-131  $\binom{131}{53}I$ ).

Complete the table.

mass number	131
number of protons	53
number of neutrons	

(1)

(b) Complete the sentence.


The number of protons in an atom is called the proton number or

the ..... number.

(1)

- (c) An atom of iodine-131 decays into an atom of xenon (Xe) by emitting a beta particle.
  - (i) The decay of iodine-131 can be represented by the equation below.

Complete the equation by writing the correct number in each of the **two** boxes.




(ii)	A sample of rainwater contaminated with iodine-131 gives a count rate of 1200 counts per second.	e.co.uk
	Calculate how many days it will take for the count rate from the sample of rainwater to fall to 75 counts per second.	
	Half-life of iodine-131 = 8 days	
	Show clearly how you work out your answer.	
	days	(2)
(iii)	If people drink water contaminated with iodine-131, the iodine-131 builds up in the thyroid gland. This continues until the thyroid is saturated with iodine-131 and cannot absorb any more. The radiation emitted from the iodine-131 could cause cancer of the thyroid.	
	In Japan, people likely to be drinking water contaminated with iodine-131 were advised to take tablets containing a non-radioactive isotope of iodine.	
	Suggest why this advice was given.	
	(Total 8 ma	(2) rks)

(1)

6

A teacher used a Geiger-Műller (GM) tube and counter to measure the *background* (a) radiation in her laboratory.

The teacher reset the counter to zero, waited one minute and then took the count reading. The teacher repeated the procedure two more times.



(i)	Background radiation can be either from natural sources or from man-made sources.
	Name one man-made source of background radiation.

(ii) The three readings taken by the teacher are given in the table.

Count after one minute
15
24
18

The readings given in the table are correct.

Why are the readings different?	

(b) Some scientists say they have found evidence to show that people living in areas of high natural background radiation are less likely to develop cancer than people living in similar areas with lower background radiation.

The evidence these scientists found does not definitely mean that the level of background radiation determines whether a person will develop cancer.

Suggest a reason why.

.....

(1)

(c) An atom of the isotope radon-222 emits an alpha particle and decays into an atom of polonium.

An alpha particle is the same as a helium nucleus. The symbol below represents an alpha particle.

4 He

(i) How many protons and how many neutrons are there in an alpha particle?


Number of protons = .....

Number of neutrons = .....


(2)

(ii) The decay of radon-222 can be represented by the equation below.

Complete the equation by writing the correct number in each of the **two** boxes.



(d) The graph shows how, in a sample of air, the number of radon-222 nuclei changes with time.



Use the graph to find the half-life of radon-222.

Show clearly on the graph how you obtain your answer.


- **7** Certain types of atom emit alpha, beta or gamma radiation. The radiation is emitted from the centre of the atom.
  - (a) What name is given to the centre of an atom?

(b) The sign below is used to warn people that a radiation source is being used in a laboratory.



	Why is it important to warn people that a radiation source is being used?	
		(1)
(c)	Before using a radiation source, a teacher asked her class whether there was any way that she could reduce the amount of radiation that the source emitted. Three students each gave an answer to the teacher.	
	Keep the source in a freezer. It will emit less radiation.  Put it in acid. It will destroy the radiation.  You can't do anything to change the amount of radiation emitted.	
	A B C	
	Which <b>one</b> of the students, <b>A</b> , <b>B</b> or <b>C</b> , is correct?	
	Write your answer in the box.	
		(1)

(d) The diagram shows the apparatus used by the teacher to demonstrate how one type of radiation is able to pass through lead.




One lead sheet, 2 mm thick, was placed between the source and the detector and a count rate was taken. Extra lead sheets were added. For each extra lead sheet, a new count rate was taken and recorded in the table.

Number of lead sheets	Count rate in counts per minute
1	226
2	220
3	210
4	190
5	185


The diagram shows how a company detects any boxes left empty by an automatic filler. (e)

When an empty box passes between the beta source and the detector, a buzzer sounds. A worker then removes the box from the conveyor belt.



(i)	Why would this system <b>not</b> work if an alpha source were used instead of the beta source?	
		(1)

The chart shows how the detector reading changes as boxes pass along the (ii) conveyor belt.




Which part of the chart, K, L, M or N, shows that an empty box is passing between the beta source and the detector?

Give a reason for your answer.	
	(2) (Total 8 marks)

Page 16 of 99

(a) The graph shows how the count rate from a sample containing the radioactive substance cobalt-60 changes with time.



(i) What is the range of the count rate shown on the graph?

From ...... counts per second to ..... counts per second.

(1)

(ii) How many years does it take for the count rate to fall from 200 counts per second to 100 counts per second?


(1)

(iii) What is the half-life of cobalt-60?

www.tutorzone.co.uk

The gamma radiation emitted from a source of cobalt-60 can be used to kill the bacteria on fresh, cooked and frozen foods. Killing the bacteria reduces the risk of food poisoning.

The diagram shows how a conveyor belt can be used to move food past a cobalt-60 source.



source has thick metal shie Which <b>one</b> of the following Draw a ring around your an	metals should be used?	?	
		?	
source has thick metal shie	siding.		
Move the cobalt-60 source	away from the conveyor		
Make the conveyor belt mo	ve more slowly.		
Increase the temperature o	f the cobalt-60 source.		
Put a tick ( $\checkmark$ ) in the box ne	ext to your answer.		
Which <b>one</b> of the following the food receives?	gives a way of increasin	ng the amount of gamma radiation	
	M	loving conveyor belt	
t	The food receives?  Put a tick ( v ) in the box ne	Which <b>one</b> of the following gives a way of increasing the food receives?  Put a tick (✓) in the box next to your answer.  Increase the temperature of the cobalt-60 source.  Wake the conveyor belt move more slowly.  Move the cobalt-60 source away from the conveyor protect people from the harmful effects of the gas	Put a tick (✓) in the box next to your answer.  ncrease the temperature of the cobalt-60 source.

www.tutorzone.co.uk
A scientist has compared the vitamin content of food exposed to gamma radiation with food that has not been exposed.

The table gives the data the scientist obtained when she tested 1 kg of cooked chicken.

Vitamin	Food not exposed to gamma radiation	Food exposed to gamma radiation
	Mass in milligrams	Mass in milligrams
B6	1.22	1.35
B12	21.00	28.00
Е	3.30	2.15
Niacin	58.00	55.50
Riboflavin	2.10	2.25

Considering only this data, which one of the following is a correct	t conclusion?
Put a tick ( $\checkmark$ ) in the box next to your answer.	
Vitamin content is not affected by gamma radiation.	
Gamma radiation completely destroys some types of vitamin.	
Exposure increased the content of some types of vitamin.	

(1) (Total 6 marks)

Food irradiation is a process that exposes food to radiation. Irradiation can be used to kill the bacteria that cause food poisoning or to slow down the ripening of fresh fruit and vegetables. Frozen foods and food inside packaging can also be irradiated.

The table gives information about five radioactive isotopes. (a)

Isotope	Half-life	Radiation emitted
Caesium-134	2.1 years	beta
Cobalt-60	5.3 years	gamma
Curium-242	160 days	alpha
Strontium-90	28 years	beta
Technetium-99	6 hours	gamma

	Whi	ch of these radioactive isotopes would be most suitable for irradiating food?	
	Expl	ain the reasons for your choice.	
			(3)
(b)		y people think that food should not be irradiated. Consumer groups have said that they worried about the nutritional value and safety of eating irradiated foods.	
	(i)	Suggest <b>one</b> reason why some people may be concerned about the safety of eating irradiated food.	
			(1)

•	scientific committees in sev ave concluded that it is sa	veral countries, including S fe to eat irradiated food.	www.tutorzor weden, Canada	ne.co
These scientil	fic committees need to be	independent from governm	nent influence.	
Suggest why.				
				(1)
irradiated food	ds.	ne vitamin content of non-ir		(1)
Vitamin	Non-irradiated food in milligrams	Irradiated food in milligrams		
B6	1.22	1.35		
B12	21.00	28.00		
E	3.30	2.15		
Niacin	58.00	55.50		
Riboflavin	2.10	2.25	_	
_	ll than non-irradiated food?	s it valid to conclude that irr	adiated food is	
				(2)

		(iv)	In a restaurant, meals with ingredients that have been irradiated must be clearly identified on the menu.	
			It is important that people eating in a restaurant are given this information.	
			Suggest why.	
				(1)
	(c)		isotope caesium-137 decays by emitting beta radiation. sium-137 has a half-life of 30 years.	
		(i)	What is a beta particle, and from which part of an atom is a beta particle emitted?	
				(1)
		(ii)	A sample containing caesium-137 has a count rate of 600 counts per minute.	
			Calculate how long it would take for the count rate from the sample to fall to 75 counts per minute.	
			Show clearly how you work out your answer.	
			Time taken = years	
			(Total 11 ma	(2) rks)
10	(a)	parti The	ns of the isotope bismuth-212 decay by emitting either an alpha particle or a beta cle. equation represents what happens when an atom of bismuth-212 decays by beta ssion into an atom of polonium-212.	
			$_{83}^{212} \longrightarrow_{84}^{212} \text{Po} + \text{beta particle}$	
		(i)	The bismuth atom and the polonium atom have the same mass number (212).	
			What is the mass number of an atom?	
				(1)

	(ii)	Beta decay does <b>not</b> cause the mass number of an atom to change.	e.co.uk
		Explain why not.	
			(2)
(b)	Whe thalli	n an atom of bismuth-212 emits an alpha particle, the atom decays into an atom of um.	
		lpha particle is the same as a helium nucleus. symbol below represents an alpha particle.	
		He 2	
	(i)	The equation below represents the alpha decay of bismuth-212.	
		Complete the equation by writing the correct number in each of the two boxes.	
		212 Bi ————————————————————————————————————	
			(2)
	(ii)	It is impossible for the alpha decay of bismuth-212 to produce the same element as the beta decay of bismuth-212.	
		Explain why.	
		(Total 7 mai	(2) rks)

11

(a) The names of the three types of nuclear radiation are given in **List A**. Some properties of these types of radiation are given in **List B**.


Draw a straight line to link each type of radiation in **List A** to its correct property in **List B**.

Draw only **three** lines.

List A Type of nuclear rad	diation	List B Property of radiation
		Has the same mass as an electron
Alpha		
		Very strongly ionising
Beta		
		Passes through 10 cm of aluminium
Gamma		
		Deflected by a magnetic field but not deflected by an electric field

(3)

www.tutorzone.co.uk The diagram shows a system used to control the thickness of cardboard as it is made.



The cardboard passes through a narrow gap between a beta radiation source and a radiation detector.

The table gives the detector readings over 1 hour.

six days

Time	Detector reading
08:00	150
08:15	148
08:30	151
08:45	101
09:00	149

(i)	Between 08:00 and 08:30, the cardboard is produced at the usual, correct thickness.
	Explain how you can tell from the detector readings that the cardboard produced at 08:45 is thicker than usual.
(ii)	Which would be the most suitable half-life for the beta source?
	Draw a ring around your answer.

six months


six years

(1)

(Total 7 marks)

(III)	alpha radiation source.	
	Why not?	
		(1)

The diagram shows a system used to control the thickness of aluminium foil as it is being rolled. A radiation source and detector are used to monitor the thickness of the foil.




(a)

Which type of source, alpha, beta or gamma, should be used in this control system?
Explain why each of the other two types of source would <b>not</b> be suitable.

(3)


The chart shows how the count rate recorded by the detector varies over a short period of time time.



Use the graph to explain how the thickness of the foil changes, and how the control system responds to this change.

(c) When first used, the radiation source contains 6 micrograms of strontium-90.

The graph shows how the mass of the strontium-90 will decrease as the nuclei decay.




The control system will continue to work with the same source until  $75\,\%$  of the original strontium-90 nuclei have decayed.

Show clearly your calculation and how you use the graph to obtain your answer.

After how many years will the source need replacing?

Number of years	=
ramber of years	
	(2)
	(Total 7 marks)

(a) The diagram represents a helium atom.



(i) Which part of the atom, K, L, M or N, is an electron?

Part	
	(1)
article?	

(ii) Which part of the atom, K, L, M or N, is the same as an alpha particle?

Part
------

(1)

(b) A radioactive source emits alpha particles.


What might this source be used for?

Put a tick ( $\checkmark$ ) in the box next to your answer.

to monitor the thickness of aluminium foli as it is made in a factory	
to make a smoke detector work	

to inject into a person as a medical tracer

(c) The graph shows how the count rate from a source of alpha radiation changes with time.



What is the count rate after 4 hours?

(a)

14

	counts per second
(1)	
(Total 4 marks)	

Carbon has three naturally occurring isotopes. The isotope, carbon-14, is radioactive.

An a	tom of carbon-14 decays by emitting a beta particle.
(i)	Complete the following sentences.

The atoms of the three carbon isotopes are the same as each other because

The atoms of the three carbon isotopes are different from each other because

(ii) What is a beta particle and from what part of an atom is it emitted?


(1)

(l=)	Www.tutorzone.co.uk
(D)	Carbon-14 is constantly being made in the atmosphere, yet for most of the last million
	years, the amount of carbon-14 in the atmosphere has not changed.

How is this possible?	
	(1)
	('')

Trees take in carbon-12 and carbon-14 from the atmosphere. After the tree dies, the (c) proportion of carbon-14 that the tree contains decreases.

Graph **A** shows the decay curve for carbon-14.




(i) Lake Cuicocha in Ecuador was formed after a volcanic eruption. Carbon taken from a tree killed by the eruption was found to have a count rate of 10.5 counts per minute.

At the time of the eruption, the count rate would have been 16 counts per minute.

Use graph **A** to find the age of Lake Cuicocha.

www.tutorzone.co.uk

(ii) Finding the age of organic matter by measuring the proportion of carbon-14 that it contains is called carbon dating. This technique relies on the ratio of carbon-14 to carbon-12 in the atmosphere remaining constant. However, this ratio is not constant so the age found by carbon dating needs to be adjusted.



Graph **B** is used to adjust the age of an object found by carbon dating. The value obtained from graph **B** will be no more than 50 years different to the true age of the object.

Use graph **B** and the information above to find the maximum age that Lake Cuicocha could be.

Show clearly how you obtain your answer.

	Maximum age of Lake Cuicocha =years
(2)	Maximum age of Lake Oulcocha =years
(Total 7 marks)	

(1)

15

Some rocks inside the Earth contain a radioactive element, uranium-238. When an atom of uranium-238 decays, it gives out an alpha particle.


(a) The following statement about alpha particles was written by a student. The statement is **not** correct.

## Alpha particles can pass through a very thin sheet of lead.

Change **one** word in the statement to make it correct.

Write down your **new** statement.

(b) The graph shows how the count rate from a sample of uranium-238 changes with time.



The graph can be used to find the half-life of uranium-238. The half-life is 4 500 million years.

(i) Draw on the graph to show how it can be used to find the half-life of uranium -238.

	(ii)	There is now half as much uranium-238 in the rocks as there was when the Earth was formed.			
		How old is the Earth?			
		Draw a ring around your answer.			
		2250 million years	4500 million years	9000 million years	(1)
	(iii)	If a sample of uranium-23 half-life in a school exper	88 were available, it would not li iment.	pe possible to measure the	(-)
		Explain why.			
					(2)
				(Total	5 marks)
<b>16</b> (a)	-	atom of uranium-238 decay	ontain uranium-238, a radioacti s, it gives out radiation and cha		n
			<b>O</b> +		
		Uranium-238	Helium nucleus	Thorium-234	
	(i)	What type of radiation is	emitted when a uranium-238 a	tom decays?	
	(ii)	From which part of a urar	nium-238 atom is the radiation	emitted?	(1)
	(11)				(4)
	(iii)	Uranium-235 is another is	sotope of uranium.		(1)
		How is an atom of uraniu	m-235 similar to an atom of ura	anium-238?	

- (b) Uranium-238 has a half-life of 4500 million years.
  - (i) When the Earth was formed, there was twice as much uranium-238 in the rocks as there is now.


What is the age of the Earth?

.....

(1)

(ii) Complete the graph to show how the number of nuclei in a sample of uranium-238 will change with time.

Initially, there were 100 000 nuclei in the sample.




(Total 6 marks)

(1)

(2)

(a) The diagram represents 3 atoms, **K**, **L** and **M**.



(i) Which **two** of the atoms are isotopes of the same element?

and	l

(ii) Give a reason why the **two** atoms that you chose in part (a)(i) are:

(1) atoms of the same element
(2) different isotopes of the same element.

.....


(b) The table gives some information about the radioactive isotope thorium-230.

mass number	230
atomic number	90

(i) How many electrons are there in an atom of thorium-230?

(1)

(ii) How many neutrons are there in an atom of thorium-230?



www.tutorzone.co.uk When a thorium-230 nucleus decays, it emits radiation and changes into radium-226. (c)


$$^{230}_{90}$$
Th  $\longrightarrow$   $^{226}_{88}$ Ra + Radiation

What type of radiation, alpha, beta or gamma, is emitted by thorium-230?

Explain the reason for your answer.

(3)(Total 8 marks)


Before using a radioactive source, a teacher asked her students to suggest safety procedures 18 that would reduce her exposure to the radiation. The students made the following



(a) Which suggestion, A, B or C, would **not** reduce the exposure of the teacher to radiation?

(1)

The diagram shows how the teacher measured the distance that the radiation traveled from the source. The count-rate at different distances from the source was measured and recorded in the table.




Distance from source to detector in cm	Count-rate in counts per minute
20	85
40	81
60	58
80	53
100	23

What type of radiation was the source emitting, alpha, beta or gamma?
Explain the reasons for your choice.

(3)

 $\begin{tabular}{ll} www.tutorzone.co.uk\\ The graphs show how two groups of scientists, {\bf A} and {\bf B}, link exposure to radiation and the\\ \end{tabular}$ (c) risk of getting cancer.



(i) Complete the following sentence using a word or phrase from the box.

	decreases	has no effect on	increases	
	Both groups of scientist	s agree that a high level	of exposure to radiation	
		the risk of	getting cancer.	(1)
i)	Use the graphs to desc the level of exposure to	<u>.                                      </u>	o groups of scientists disa	
				(2) (Total 7 marks)

(1)

Most elements have some isotopes which are radioactive.


(a) What is meant by the terms:

(i) isotopes


(ii) radioactive?

(1)

(b) The graph shows how the number of nuclei in a sample of the radioactive isotope plutonium-238 changes with time.



Use the graph to find the half-life of plutonium-238.

Show clearly on the graph how you obtain your answer.

Half-life = ..... years

(2)

(c)	The	Cassini spacecraft launched in 1997 took seven years to reach Saturn.	rzone.co
		electricity to power the instruments on board the spacecraft is generated using the produced from the decay of plutonium-238.	
	(i)	Plutonium-238 decays by emitting alpha particles.	
		What is an alpha particle?	
	(ii)	During the 11 years that Cassini will orbit Saturn, the output from the generators will decrease.	(1)
		Explain why.	
			(2)
(d)	Pluto hum	onium-238 is highly dangerous. A tiny amount taken into the body is enough to kill a lan.	
	(i)	Plutonium-238 is unlikely to cause any harm if it is outside the body but is likely to ki if it is inside the body.	II
		Explain why.	
			(2)
	(ii)	In 1964, a satellite powered by plutonium-238 was destroyed, causing the release of radioactive material into the atmosphere.	
		Suggest why some environmental groups protested about the launch of Cassini	

(1) (Total 10 marks)

20

(a) Complete the following table for an atom of uranium-238 (  $\frac{238}{92}$  U)

mass number	238
number of protons	92
number of neutrons	

(1)

(b) Complete the following sentence.

The name given to the number of protons in an atom is the proton number or the

(1)

(c) An atom of uranium-238 (  $\frac{238}{92}$  U) decays to form an atom of thorium-234 (  $\frac{234}{90}$  Th).

(i) What type of radiation, alpha, beta or gamma, is emitted by uranium-238?

(1)


(ii) Why does an atom that decays by emitting alpha or beta radiation become an atom of a different element?

(Total 4 marks)

Total I manto,

**21** In

The diagram shows a helium atom.



(a) (i) Use the words in the box to label the diagram.

(2)


(ii) An alpha particle is the same as the nucleus of a helium atom.

How is an alpha particle different from a helium atom?

.....

(1)

(b) The graph shows how the count rate from a sample of radioactive sodium-24 changes with time.



(i) How many hours does it take for the count rate to fall from 100 counts per second to 50 counts per second?

Time = ..... hours

(1)

(ii) What is the half-life of sodium-24?


Half-life = ..... hours

(1)

(c)	A sn	noke detector contains a small amount of americium-241.	.00.0			
		Americium-241 is a radioactive substance which emits alpha particles. It has a half-life of 432 years.				
	(i)	Which <b>one</b> of the following statements gives a reason why the americium-241 inside the smoke detector will <b>not</b> need replacing?				
		Put a tick (✓) in the box next to your answer.				
		The alpha particles have a low energy.				
		People replace smoke detectors every few years.				
		Americium-241 has a long half-life.	(1)			
	(ii)	The diagram shows the label on the back of the smoke detector.				
		Red Tive Material				
		Why do people need to know that the smoke detector contains a radioactive material?				
		(Total 7 mark	(1) ks)			

(i)

The diagram shows the parts of a smoke detector. The radioactive source emits alpha (a) particles.




The alpha particles ionise the air inside the sensor which causes a small electric current. Any smoke getting into the sensor changes the current. The change in current sets the alarm off.

The smoke detector would **not** work if a radioactive source that emitted only gamma

	rays was used.	
	Why not?	
		(1)
(ii)	Curium-242 is a radioactive isotope with a half-life of 160 days. It emits alpha particles.	
	Why is curium-242 <b>not</b> suitable for use inside smoke detectors?	
		(1)
(iii)	Curium-242 and curium-244 are two of the isotopes of the element curium.	,
	How is an atom of curium-242 different from an atom of curium-244?	
		(1)
		\ · /

(b) Sections of steel are often joined by welding them together. The diagram shows how a radioactive source can be used to check for tiny cracks in the weld.




Cracks in the weld will be shown up on the photographic film below the thick steel plate.

(i)	Which type of source, alpha, beta or gamma, should be used to check the weld?			
		(1)		
(ii)	Give a reason why the other two types of source <b>cannot</b> be used.			
		(1)		

(c) The diagram shows a map of a river and its estuary.

Environmental scientists have found that the water flowing into one part of the river estuary is polluted. To find where the pollution is coming from, the scientists use a radioactive isotope, gold-198.



(i)	Explai	n how th	e gold-198 is use	d to find whe	ere the pollu	ution is cor	ning from	www.tutorzone.co
								(2)
(ii)	The gr	aph sho	vs how the count	rate from a	sample of g	jold-198 cl	nanges w	vith time.
		550						
		500						
		450	N. N. H.					
		400						
Coun	t rate	350						
in co		300						
× 10	000	250						
		200						
		150				<b>**</b>		
		100						

Use the graph to calculate the half-life of gold-198.

Show clearly on the graph how you obtain your answer.


Time in days

Half-life = ..... days

(2) (Total 9 marks)

A radioactive source emits alpha  $(\alpha)$ , beta  $(\beta)$  and gamma  $(\gamma)$  radiation. The diagram shows what happens to the radiation as it passes between two charged metal plates.

## Diagram 1



Which line **P**, **Q** or **R** shows the path taken by: (a)

> (i) alpha radiation


> > (1)

(ii) gamma radiation?


(1)

(b) The diagram shows three different boxes and three radioactive sources. Each source emits only one type of radiation and is stored in a different box. The box reduces the amount of radiation getting into the air.













Gamma source





Draw three lines to show which source should be stored in which box so that the minimum amount of radiation gets into the air.

(2)

The graphs show how the count rates from three different radioactive sources,  $\mathbf{J}$ ,  $\mathbf{K}$ , and  $\mathbf{L}$ , change with time. 100 100 100 80 80 Count Count Count rate in 60 rate in 60 rate in 60 counts counts counts per 40 per 40 per 40 second second second 20 20 20 0 10 15 20 10 15 15 20 0 10 Time in hours Time in hours Time in hours J K L (i) Which source, J, K, or L, has the highest count rate after 24 hours? (1) (ii) For source **L**, what is the count rate after 5 hours? ..... counts per second (1) (iii) Which source, J, K, or L, has the longest half-life? ..... (1) A radioactive source has a half-life of 6 hours. (iv) What might this source be used for? Put a tick  $(\checkmark)$  in the box next to your choice.

To monitor the thickness of paper as it is made in a factory


To inject into a person as a medical tracer

To make a smoke alarm work

(1) (Total 8 marks)

(a)	A ra	adioactive source emits alpha $(\alpha)$ , beta $(\beta)$ and gamma $(\gamma)$ radiation.	
	(i)	Which two types of radiation will pass through a sheet of card?	
			(1)
	(ii)	Which <b>two</b> types of radiation would be deflected by an electric field?	
	(iii)	Which type of radiation has the greatest range in air?	(1)
	, ,		(1)
(b)		udent suggests that the radioactive source should be stored in a freezer at – 20 student thinks that this would reduce the radiation emitted from the source.	
		gest why the student is wrong.	
			(1)
(c)	Pho	sphorus-32 is a radioactive isotope that emits beta radiation.	(1)
	(i)	How is an atom of phosphorus-32 different from an atom of the stable isotope phosphorus-31?	
			(1)

www.tutorzone.co.uk
The graph shows how the count rate of a sample of phosphorus-32 changes with (ii) time.




Use the graph to calculate the half-life of phosphorus-32.

Show clearly how you used the graph to obtain your answer.

Half-life = ..... days

(2)


Plants use phosphorus compounds to grow. Watering the root system of a plant with (iii) a solution containing a phosphorus-32 compound can help scientists to understand the growth process.



Explain why phosphorus-32 is suitable for use as a tracer in this situation.	
	(2) (Total 9 marks)

(a) The names of three types of nuclear radiation are given in **List A**. Some properties of these three types of radiation are given in **List B**.

Draw a straight line to link each type of radiation in **List A** to its correct property in **List B**. Draw only three lines.




One of the substances in the table is used as a radioactive tracer. A hospital patient breathes in air containing the tracer. The radiation given out is measured by a doctor using a detector outside the patient's body.

Substance	Radiation given out	Solid, liquid or gas
х	alpha	gas
Υ	gamma	gas
Z	gamma	solid

	Which one of the substances, A, Y of Z, should be used as the tracer?	
	Give <b>two</b> reasons for your answer.	
	1	
	2	
		(3)
(d)	Radiation can also be used to kill the bacteria on fresh food.	
	Give <b>one</b> reason why farmers, shop owners or consumers may want food to be treated with radiation.	
	(Total 8 ma	(1) arks)

Radon is a radioactive element. The graph shows how the number of radon atoms in a sample of air changes with time.




(i) How long did it take the number of radon atoms in the sample of air to fall from 1000 to 500?

(ii) How long is the half-life of radon?

(iii) Complete this sentence by crossing out the **two** lines in the box that are wrong.

As a radioactive material gets older, it emits less a constant level of more radiation per second.

(1) (Total 3 marks) The detector and counter are used in an experiment to show that a radioactive source gives out alpha and beta radiation only.



Two different types of absorber are placed one at a time between the detector and the source. For each absorber, a count is taken over ten minutes and the average number of counts per second worked out. The results are shown in the table.

Explain how these results show that alpha and beta radiation is being given out, but gamma

Absorber used	Average counts per second
No absorber	33
Card 1 mm thick	20
Metal 3 mm thick	2


radiation is <b>not</b> being given out.	
	(Total 3 marks)

(a) The table gives information about six radioactive isotopes.

Isotope	Type of radiation emitted	Half-life
hydrogen-3	beta particle	12 years
iridium-192	gamma ray	74 days
polonium-210	alpha particle	138 days
polonium-213	alpha particle	less than 1 second
technetium-99	gamma ray	6 days
uranium-239	beta particle	24 minutes

(i)	What is an alpha particle?	
		(1)
(ii)	Two isotopes of polonium are given in the table. How do the nuclei of these two isotopes differ?	
		(1)
(iii)	A doctor needs to monitor the blood flow through a patient's heart. The doctor injects a radioactive isotope into the patient's bloodstream. The radiation emitted by the isotope is then detected outside the body.	
	Which <b>one</b> of the isotopes in the table would the doctor inject into the bloodstream?	
	Explain the reasons for your choice.	
		(3)

(b) Igneous rock contains uranium-238 which eventually changes to the stable isotope lead-206. The graph shows how the percentage of uranium-238 nuclei present in an igneous rock changes with time.



A rock sample is found to have seven atoms of uranium-238 for every three atoms of lead-206. Use the graph to estimate the age of the rock. Show clearly how you obtain your answer.

 	 	 • • • • • •

(Total 7 marks)


A smoke detector fitted inside a house contains a radioactive source, americium 241.

(a) Complete the following table of information for an atom of americium 241.

29

Number ofneutrons	146
Number ofprotons	95
Number ofelectrons	

(b)	The diagram shows that the radiation given out by americium 241 does not go	www.tutorzone.co.ul through
	paper.	



Radiation / Thin paper	
Which type of radiation, alpha ( $\alpha$ ), beta ( $\beta$ ), or gamma ( $\gamma$ ) is given out by americium 241?	
Explain why the radiation given out by the americium 241 is unlikely to do any harm to people living in the house.	(1)
Complete the sentence by choosing an answer from the box.	(2)
less than more than the same as	
After many years the radiation emitted by americium 241 will be	
when the smoke detector was new.	(4)
(Total s	(1) 5 marks)
eta particle is a high-energy electron.	
Which part of an atom emits a beta particle?	
	(1)
	Which type of radiation, alpha (α), beta (β), or gamma (γ) is given out by americium 241?  Explain why the radiation given out by the americium 241 is unlikely to do any harm to people living in the house.  Complete the sentence by choosing an answer from the box.  Iess than more than the same as  After many years the radiation emitted by americium 241 will be

30


(ii) How does the composition of an atom change when it emits a beta particle?

.....

(1) (Total 2 marks)

31

(a) Radiation can cause cancer. The graph shows that the risk of cancer depends on the radiation dose a person is exposed to.



Complete the following sentence.

The ...... the dose of radiation a person gets, the greater the risk of cancer.

(1)

(b) A worker in a nuclear power station wears a special badge (diagram 1). Diagram 2 shows what is inside the badge. When the film inside the badge is developed, it will be dark in the places where it has absorbed radiation.




Diagram 1

Diagram 2

.....

Give a reason for your answer.

.....

(Total 3 marks)


The table shows how the count rate from a radioactive substance changes in 10 days.

Time in days	0	2	4	6	8	10
Count rate in countsper minute	880	555	350	220	140	90

(a) Draw a graph of count rate against time.

**32** 

The first two points have been plotted for you.



(3)

.uk

(b)	(i)	Use your graph to find o	_	es for the count rate to fa	www.tutorz all from 880 counts	one.co.
			Ti	me =	days	(1)
	(ii)	What is the half-life of the	nis substance?			
			Half-	life =	days	(1)
(c)		table gives the half-life aropes.	nd type of radiation	n given out by four differ	ent radioactive	, ,
		Radioactive isotope	Half-life in days	Radiation given out		
		bismuth-210	5.0	beta		
		polonium-210	138.0	alpha and gamma		

3.8

24.1

Some samples of each isotope have the same count rate today. Which sample with the lowest count rate one month from today?	II nave
Give a reason for your answer.	
	(2)
	(Total 7 marks)

alpha

beta and gamma

Read the information in the box and then answer the questions.

33

radon-222

thorium-234

Igneous rocks contain potassium-40. This is a radioactive isotope. It has a half-life of 1300 million years.

Potassium-40 decays into argon-40 which is stable.

Argon escapes from molten rock. Any argon found in an igneous rock must have been produced since the rock solidified.

A sample of an igneous rock has one atom of potassium-40 for every three atoms of argon-40.

(i) What fraction of the potassium-40 has not yet decayed?

(1)

(ii) Calculate the age of the rock.


34

.....

Age of rock = ..... million years

(1) (Total 2 marks)

(a) The graph shows how a sample of barium-143, a radioactive *isotope* with a short *half-life*, decays with time.



www.tutorzone.co.uk (i) What is meant by the term *isotope?* (1) (ii) What is meant by the term *half-life*? (1) (iii) Use the graph to find the half-life of barium-143. Half-life = ..... seconds (1) Humans take in the radioactive isotope carbon-14 from their food. After their death, the (b) proportion of carbon-14 in their bones can be used to tell how long it is since they died. Carbon-14 has a half-life of 5700 years. A bone in a living human contains 80 units of carbon-14. An identical bone taken from (i) a skeleton found in an ancient burial ground contains 5 units of carbon-14. Calculate the age of the skeleton. Show clearly how you work out your answer. Age of skeleton = ..... years (2) Why is carbon-14 unsuitable for dating a skeleton believed to be about 150 years (ii) old?

(1)

(c) The increased industrial use of radioactive materials is leading to increased amounts of radioactive waste. Some people suggest that radioactive liquid waste can be mixed with water and then safely dumped at sea. Do you agree with this suggestion? Explain the reason for your answer.


(3) (Total 9 marks)

(a) What is a beta (β) particle?

35

(1)

(b) Plants absorb carbon-14 from the atmosphere. The graph shows the decay curve for 1 g of carbon-14 taken from a flax plant.



Use the graph to find the half-life of carbon-l4. You should show clearly on your graph how you obtain your answer.

tait-lite =	years.
	(2)

(c) Linen is a cloth made from the flax plant. A recent exhibition included part of a linen shirt, believed to have belonged to St. Thomas à Becket, who died in 1162. Extracting carbon-14 from the cloth would allow the age of the shirt to be verified.

If 1 g of carbon-14 extracted from the cloth were to give 870 counts in 1 hour, would it be possible for the shirt to have once belonged to St. Thomas à Becket? You must show clearly the steps used and reason for your decision.


Page 66 of 99

(3)


(Total 6 marks)

(a) The table gives information about five radioactive isotopes.

Isotope	Type of radiation emitted	Half-life
Californium-241	alpha (α)	4 minutes
Cobalt-60	gamma (γ)	5 years
Hydrogen-3	beta (β)	12 years
Strontium-90	beta (β)	28 years
Technetium-99	gamma (γ)	6 hours

(1)	What is an alpha (α) particle?	
		(1)
(ii)	What is meant by the term half-life?	
		(1)
(iii)	Which <b>one</b> of the isotopes could be used as a tracer in medicine? Explain the reason for your choice.	
		(3)

www.tutorzone.co.uk
The increased use of radioactive isotopes is leading to an increase in the amount of
radioactive waste. One method for storing the waste. (b) radioactive waste. One method for storing the waste is to seal it in containers which are then placed deep underground.



they live. Explain why.	
	(2)
	(3) (Total 8 marks)

38

(a) The diagram shows a hazard sign.



What type of hazard does this sign warn you about?


.....(1)

(b) The names of three types of radiation are given in the box.

alpha (α)	beta (β)	gamma (γ)	
-----------	----------	-----------	--

Complete each sentence by choosing the correct type of radiation from those given in the box. Each type of radiation should be used once or not at all.

(a) A radiation detector and counter were used to detect and measure the radiation emitted from a weak source. The graph shows how the number of counts recorded in one minute changed with time.




www.tutorzone.co.uk Even though the readings from the counter were accurately recorded, not all the points fit the smooth curve. What does this tell us about the process of radioactive decay? (1) (ii) After ten minutes the number of counts recorded each minute is almost constant. Explain why. (2) (b) The radioactive isotope sodium-24 injected into the bloodstream can be used to trace blood flow to the heart. Sodium-24 emits both beta particles and gamma rays. (i) What is a beta particle? (1) (ii) What is a gamma ray? (1) (iii) The count rate from a solution containing sodium-24 decreases from 584 counts per minute to 73 counts per minute in 45 hours. Calculate the half-life of sodium-2.2. Show clearly how you work out your answer. Half-life = ..... hours

(3)

		(iv)	Give <b>one</b> advantage of using sodium-24 to trace blood flow compared to usotope with a half-life of:	www.tutorzone.using an	co.uk
			[A] ten years;		
			[B] ten seconds.	(	1)
				(Total 10 mark	1) s)
39	(a)	only You	o sources of radiation look identical. One source emits only alpha radiation, to beta radiation. Describe <b>one</b> way to find out which source emits the alpha can assume a radiation detector and counter are available. You may wish to help with your answer.	radiation.	
				(	3)

www.tutorzone.co.uk
The diagram shows a beta radiation source and detector used to measure the thickness of (b) cardboard as it is made. The table gives the detected count rate at different times.



Time	Count rate in counts/minute
09:00	120
09:30	122
10:00	119
10:30	165
11:00	118

1)	thickness. Give a reason for the small variation in count rate.	
		(1)
ii)	What can you say about the thickness of the cardboard being made at 10:30?	(1)
	Explain the reason for your answer.	
		(3)

Radon is a radioactive gas. Radon makes a major contribution to background radiation levels. Radon atoms decay by the emission of *alpha particles*.


(a) (i) What is an alpha particle?

(1)

(ii) From which part of the radon atom does the alpha particle come?

(1)

(b) (i) A sample of air contains 40 000 radon atoms. The half-life of radon is four days. Draw a graph to show how the number of radon atoms present in a sample of air will change over a period of 12 days.



Time in days


(3)

(Total 8 marks)

uk

	(ii)	After 20 days, how many of the radon atoms from the original sample of air will have decayed? Show clearly how you work out your answer.	1e.co.ı
		Number of radon atoms decayed =	(3)
(c)	Fairl	y constant concentrations of radon gas have been found in some deep mine shafts.	(0)
	(i)	Suggest why the concentration of radon gas remains fairly constant although the radon gas decays.	
			(1)
	(ii)	Explain why the long term exposure to large concentrations of radon gas could be a danger to health.	
			(2)
		(Total 11 ma	arks)

The diagram below shows a method of controlling the thickness of paper produced at a paper mill. A radioactive source which emits beta radiation is placed on one side of the paper and a radiation detector is placed on the other.



41

(a)	How	www.tutorzo	one.co.uk
			(1)
(b)	Expl	ain, as fully as you can:	
	(i)	why a radioactive source which emits alpha $(\alpha)$ radiation could $\boldsymbol{not}$ be used for this application.	
			(1)
	(ii)	why a radioactive source which emits gamma ( $\gamma$ ) radiation could <b>not</b> be used for this application.	
	(iii)	why a radioactive source which emits beta $(\beta)$ radiation <b>can</b> be used for this application.	(1)
			(2)

(c)	www.tutorzone.co.uk Americium-241 is a radioisotope used in smoke detectors. It has a proton number of 95 and a mass number of 241.
	How long would it take the americium-241 in a smoke detector to decrease to one eighth of


its original number of radioactive atoms?
Answer =

(3)(Total 8 marks)

(a)

Tritium  $({}^3_1 \text{H}\,)$  is an isotope of hydrogen. Tritium has a proton number of 1 and a mass number of 3.

The diagram below shows a simple model of a tritium atom. Complete the diagram by (i) adding the names of the particles indicated by the labels.




(4)

		( - /
(ii)	Explain how the nucleus of an ordinary hydrogen atom is different from the nucleus of	
	a tritium atom. Ordinary hydrogen atoms ( $^1_1 H$ ) have a mass number of 1.	
		(2)
(iii)	Tritium is a radioactive substance which emits beta (β) radiation.	(-)
(111)	Why do the atoms of some substances give out radiation?	

(2)

www.tutorzone.co.uk Tritium is one of the elements found in the waste material of the nuclear power industry. (b) The diagram below shows a worker behind a protective screen. The container holds a mixture of different waste materials which emit alpha ( $\alpha$ ), beta ( $\beta$ ) and gamma ( $\gamma$ ) radiation.



Suggest a suitable material for the protective screen. The material should prevent radiation from the container reaching the worker. Explain your answer.	
(Total 10 ma	(2) arks)

(a)	Complete the tabl	le about atomic partic	les

ATOMIC PARTICLE	RELATIVE MASS	RELATIVE CHARGE
proton		+1
neutron	1	0
electron	negligible	


(2)

(b)	Use	the Data Sheet to help you to answer some parts of this question.	tutorzone.co.l			
	Rea	Read the following passage about potassium.				
		assium is a metallic element in Group 1 of the Periodic Table. as a proton (atomic) number of 19.				
	lts m	nost common isotope is potassium-39, ( $^{40}_{19}$ K ).				
	Anot	ther isotope, potassium-40, ( $\frac{40}{19}$ K), is a radioisotope.				
	(i)	State the number of protons, neutrons and electrons in potassium-39.				
		Number of protons				
		Number of neutrons				
		Number of electrons	(2)			
	(ii)	Explain why potassium-40 has a different mass number from potassium-39.				
			(1)			
	(iii)	What is meant by a radioisotope?	(-)			
	(iv)	Atoms of potassium-40 change into atoms of a different element. This element has proton (atomic) number of 20 and a mass number of 40.	(1) nas a			
		Name, or give the symbol of, this new element.				
			(1)			
	(v)	Explain in terms of atomic structure, why potassium-39 and potassium-40 have to same chemical reactions.	he			
			(1)			
(c)	(i)	Name a suitable detector that could be used to show that potassium-40 gives our adiation.	t			
			(1)			

www.tutorzone.co.uk

(ii) Name a disease which can be caused by too much exposure to a radioactive substance such as potassium-40.	J.,
(1) (Total 10 marks	
<ul> <li>(a) A radioactive isotope has a half-life of 10 minutes.         At the start of an experiment, the activity of a sample of this isotope was 800 counts per second after allowing for background radiation.     </li> <li>Calculate how long it would be before the activity fell from 800 counts per second to 200 counts per second.</li> </ul>	
Time min. (2)	)
<ul> <li>(b) A physicist investigates a solid radioactive material. It emits alpha particles, beta particles and gamma rays.         The physicist does not touch the material.     </li> <li>Explain why the alpha particles are less dangerous than the beta particles and gamma rays.</li> </ul>	

The diagram shows a film badge worn by people who work with radioactive materials. The badge has been opened. The badge is used to measure the amount of radiation to which the workers have been exposed.




		A window B
(a)		detector is a piece of photographic film wrapped in paper inside part <b>B</b> of the badge. <b>A</b> has "windows" as shown.
	Com	plete the sentences below.
	Whe	en the badge is closed
	(i)	radiation and radiation can pass through the open
		window and affect the film.
	(ii)	Most of the radiation will pass through the lead window and
		affect the film.
(b)	Othe	er detectors of radiation use a gas which is ionised by the radiation.
	(i)	Explain what is meant by ionised.
	(ii)	Write down <b>one</b> use of ionising radiation.

(1)

(c)		www.tutorzone.co.u nium-238 has a very long half-life. It decays via a series of short-lived radioisotopes to uce the stable isotope lead-204.	лk
	Expl	ain, in detail, what is meant by:	
	(i)	half-life,	
	(ii)	radioisotopes. (1)	
		(2)	
(d)	The	relative proportions of uranium-238 and lead-204 in a sample of igneous rock can be	
( )	usec	I to date the rock. ck sample contains three times as many lead atoms as uranium atoms.	
	(i)	What fraction of the original uranium is left in the rock?	
	(.)	(Assume that there was no lead in the original rock.)	
		( cooms that the rate has no rought and ongine rought)	
		(1)	
	(ii)	The half-life of uranium-238 is 4500 million years.	
		Calculate the age of the rock.	
		Age million years (2)	
		(Total 10 marks)	

The diagram shows a badge worn by a worker at a nuclear power station.

Part of the outer black paper has been removed so that you can see the inside of the badge.



Scientists examined the worker's badge at the end of a day's work.

They found that the top part of the badge had been affected by radiation, but the bottom half had not.

	(Total 2 marks)
What type of radiation had the worker been exposed to? Explain the reasons for your ans	SWC1.
What tung of radiation had the worker heen evinced to 7 Evniain the reacons for volir and	ς\MΔr

The diagram shows how the thickness of aluminium foil is controlled. The thicker the aluminium foil, the more radiation it absorbs.

Pressure control Roller	Detector	Aluminium foil
	Beta radiation source	

(a)	The	designers used a beta radiation source for this control system.	
	(i)	Why would an alpha radiation source be unsuitable in this control system?	
			(1)
	(ii)	Why would a gamma radiation source be unsuitable in this control system?	
			(1)
(b)	The	substance used in the beta radiation source is radioactive.	
	(i)	Why are some atoms radioactive?	
			(1)
	(ii)	Explain why radiation is dangerous to humans.	
			(2) (Total 5 marks)


(a)	(i)	Describe the structure of alpha particles.	www.tutorzone.co.uk
			(2)
	(ii)	What are beta particles?	

(b) Describe how beta radiation is produced by a radioactive isotope.

(1) (Total 4 marks)

49 lodine-131 (131) is a radioactive isotope used in medicine.

The graph shows how the count rate of a sample of iodine-131 changed over 24 days.



	(i)		the graph to calculate the half-life of iodine-131. To obtain full marks you sh ly how you work out your answer.	www.tutorzone.co.uk ould show
			Half-life days	
	(ii)	lodir	ne-131 is used to destroy cancer cells in the human thyroid gland.	(2)
		Expl	lain why the length of the half-life of iodine-131 is important in this use.	
				(2)
				(2) (Total 4 marks)
	(-)	<i>(</i> ')		
50	(a)	(i)	Describe the structure of alpha particles.	
				(2)
		(ii)	What are beta particles?	
				(1)

A teacher reach week,					tion fro	m a rad	lioacti	ve soul	rce, dur	ing the	same	lesson
The results	are show	wn on th	ne grap	h.								
Radiation in counts per minute	70 60 50 40 30 10											
	Ö	1		2		3		4	5		б	
How long d			out you	ur ansv	fall froi ver.		ounts p		ute to h			?

52	$^{99}_{43} \text{Tc}$ (technetium) is produced by the radioactive decay of $^{99}_{42} \text{Mb}$ (molybdenum).	WWW.dialog.com
	What change occurs in the nucleus of a molybdenum atom when this happens?	
		(Total 1 mark)

The isotope of sodium with a mass number of 24 is radioactive. The following data were obtained in an experiment to find the half-life of sodium-24.

53

Time in hours	Count rate in counts per minute
0	1600
10	1000
20	600
30	400
40	300
50	150
60	100

Draw a graph of the results and find the half-life for the isotope. On the graph show how you obtain the half-life.


	+++	++++		+++	++++	+++			нн	+++	+++		+++	+++	++	+++	++-		++	+++	++	-	н
	+++	+++	+++	+++	++++	+++			+++		+++	-	+	+	++	+++	++	++	++	+++	++	++-	+
	+++	++++	111	+++	++++	+++	-		нн		+	_	+	+++	++	+++	++	+++	++	+++	++	-	Н
			111	$^{\rm H}$	-	***	$\mathbf{H}$		ш		1111			111		$^{\rm HI}$	+	ш	**	+	11		ш
					$\overline{}$	$\mathbf{H}$										$^{\rm H}$		$\Box$	**	$\mathbf{H}$			$\Box$
																			$^{+}$				$\Box$
																			П	ш			
															ш	ш	ш		ш	ш			
																			ш				
						-			$\perp$		$\perp$		$\perp$	$\perp$	-					$\perp$	-		$\perp$
					+				$\perp$	+	$\perp$			$\perp$	$\perp$	$\Box$	+	$\sqcup$	-	$\perp$	+		$\perp$
	+++	++++			+++		$\perp$		ш	+++	+++	-	$\perp$	+	11	+++	++	$\vdash$	++	+++		Н-	4
	+++	++++		+++	++++		+		+++		+++	-	+++	+++	-	+++	-	$\vdash$	++	+++		Н-	
	+++	+++		+++	++++	+++	+++		нн	+++	+++		+++	+++	-	+++	++	+++	++	+++	++-	-	+
	+++	++++		+++		+++		+++	нн		+++	+	+++	+++	++	+++	++-	+++	++	+++	++	-	ш
	+++	++++	+++	+++	++++	+++			нн		+++	-	+++		++	+++	++-	ш	++	+++	++	-	н
	+++	++++	+++	+++	++++	+++	+++		нн	+++	+++	+	+++	+	++	+++	++	+++	++	+++	++	-	Н
		1111	+++		++++	+++	+++	-	нн	+++	+++		+++	-	++	111	++	H	++	111	++		Н
	-	++++	+++	+++	++++	+++	-		нн	***	+++	-	-	111	++	+++	++	++	++	111	++	-	Н
	$\overline{}$	++++	+++	+++	1111	+	$\mathbf{H}$		н	+++	$^{\rm HI}$		+	111	-	$^{++}$	++	н	++	111	11	+	Н
			111		1111	111			-		$\mathbf{H}$		-	111	11	111	11	$\Box$	11	$^{\rm H}$	11		ш
																$\Box$							П
										$\perp$									ш				
					$\perp$										11		$\perp$		$\perp$		$\perp$		
					$\perp$	1									11	$\Box$	-	$\vdash$	$\perp$		$\perp$		$\perp$
		$\perp$	111		$\perp$		$\perp$		$\perp$	+++	+++			$\perp$	1	$\perp$	++	$\square$	11	$\square$	-	1	44
	+++	+++			+++						+++		+++	+++	++	+++	++-	$\vdash$	++	+++	++	-	ш
	+++	+++	111		++++	111			ш		+++	-	+++	+	++	+++	++-	H	++	+++	++	-	ш
	+++		+++						нн		+++		+++	+++	-	+++	++-	H	++	+++	++	-	ш
	+++	++++		+++	++++	+++			нн	+++	+++	-	+++		++	+++	++-	H	++	111	++	-	ш
	+++	1111	+++	+++	++++	+++	+		нн		+++	-	+++	+++	++	++	++	+++	++	+++	++	-	Н
	+++	++++	111		++++	+++	+		нн	+++	+++	-	+++	+++	++	+++	++	H	++	111	++		Н
	+++	++++	111	HH	1111	111	+	-	нн	-	1111			+++	++	111	++	+	++	111	++		Н
		++++	111	+	1111	+	$^{\rm H}$	-	нн	-	$^{\rm HH}$		+	111	-	$^{\rm HI}$	++	$^{\rm H}$	++	111	++		Н
Count rate			111		1111					111			111	111	11		11		11		11		т
																П							П
in																$\Box$							
***																							
counts per														1.1.1	11				$\Pi$				
counts per											$\perp$		$\perp$	$\perp$	11						-		$\perp$
minute					$\perp$				ш	111				$\perp$	$\perp$				11		1		ш
minute		1111	111		+	111			+++		$\perp$		$\perp$	$\perp$	1	H	11	$\perp$	11	$\square$	-	1	ш
	+++	1111	+++	+++	+++				+++		1111		+++		++	H	++	+++	++	111	++		ш
	+++	++++	+++	+++	++++	+++	+++	+++	нн		HH	+	+++	+++	++	+++	++-	HH	++	+++	++	-	н
	+++	++++	+++		++++	+++	-		нн	+++	+++	-	+++	+++	++	+++	++	+++	++	111	++		н
	+++	++++	+++	+++	++++	+++	+++	+++	нн	+++	+++	+	+++	+++	++	+++	+	+++	++	+++	++	++	Н
	+++	++++	111	+++	++++	+++	+		нн		+++		+++	+++	++	+++	++	111	++	111	++	-	+
									нн		$^{\rm H}$			111				$\mathbf{H}$	**	+			ш
																			11				ш
																			П	ш			
																ш			$\perp$	ш			
		++++		+++	+++		+		ш		+++	-	+++	+	-	$\vdash$	-	$\vdash$	++	ш	-	Н.	-
	+++	+++		+++	++++		+++		ш	+++	+++	-	+++	+++	++	+++	++-	$\vdash$	++	+++	++	-	
	$\rightarrow$		+++	+++		+++			нн	+++		-	+++	+++	++		++	$\mapsto$	++	+++	++	-	ж
		++++			++++	+++	+++		нн		+++		+++	+++	++	+++	++	+++	++	+++	++	-	+
	+++	++++	+++	+++	++++	+++	+++		нн		+++		+++		++	+++	++-	++	++	+++	++	-	н
	+++	++++	111		++++	+++	-		нн		+++	+	+++	+++	11	111	++	++	++	+++	++	-	Н
	+++	1111	111			+++	-		нн					-	-	111	++	111	++	+++	++	-	Н
					1111	111			нн	111				111		+	11	$\Box$	**	$\mathbf{H}$	11		ш
		$\Box$			1111					-				111	1		1		11	$\Box$	-	1	
		+	111		1111	111			+++		+++	-	$\perp$	$\perp$	11	$\Box$	1	11	11	$\Pi$	1		+
		++++	111	+++	++++				+++		+++		+++	+	++	111	-	+	++	+++	++	-	+
	+++		111	+++	++++	+++			+++	+++	+++	-	+++	+++	++	+++	++	+++	++	+++	++	-	+
	+++	++++	+++	+++	++++	+++		+++	+++	+++	+++	-	+++	+++	++	+++	++	+++	++	+++	++	-	+
	+++	++++	111	+++	++++	111	1111			+++	1111	-		+++	++	+++	++	+++	++	+++	++	++-	Н
	+++	++++	+++	1111	++++	+++	-			+++	1111	-		+++	++	+++	++	+++	++	+++	++	1	+
		1111	111	+++		111			1111	+++	+++		111	111	11	111	11		11	111	+		+
																_				_		_	_

Time in hours

Half-life = ..... hours

(4)

www.tutorzone.co.uk Sodium-24 decays by beta emission. The G.M. tube used in the experiment is shown in the (b) diagram. Each beta particle which gets through the glass causes a tiny electric current to pass in the circuit connected to the counter.



	(i)	Why must the glass wall of the G.M. tube be very thin?	
			(1)
	(ii)	Why is this type of arrangement of no use if the radioactive decay is by alpha emission?	
			(1)
c)	patie	um chloride solution is known as saline. It is the liquid used in 'drips' for seriously-ill ents. Radioactive sodium chloride, containing the isotope sodium-24, can be used as a er to follow the movement of sodium ions through living organisms.	
		one advantage of using a sodium isotope with a half-life of a few hours compared to g an isotope with a half-life of:	
	(i)	five years;	
			(1)
	(ii)	five seconds.	
		/T-4-10	(1)
		(Total 8 ma	rks)

People who work in places where radiation is present, for example in X-ray departments in hospitals, have to wear a "film badge". These badges are sent away regularly to check on the amount of radiation to which the person has been exposed. Simply described, the badge is some photographic film in a suitable holder.



(a)	(i)	Why is the "film badge" of little use in detecting alpha particles?	
			(1)
	(ii)	How does the "film badge" show radiation has reached it?	
			(1)
(b)	Radi	oactivity can cause harm. It also has a number of valuable uses.	
	(i)	How can radioactivity harm our bodies?	
			(1)
	(ii)	Give <b>two</b> medical uses of radioactive isotopes.	
		1	
		2	(2)

(c)	A ra	dioactive isotope of lead has a half-life of 10.6 hours.	
	A sr	nall sample of lead containing this isotope has a count rate of 8000 counts per minute.	
	How	long will it be before the count rate is 1000 counts per minute?	
		Time = hours	
		(Total 7 ma	(2) arks)
		· ·	,
clos	e toge	spark counter can be used to detect charged particles. It is made by having two wires other with a large voltage across them. When a charged particle passes through the seen the wires a spark is seen.	
(a)	Give	e the names and symbols of <b>two</b> particles which will cause a spark.	
	(i)	Name Symbol	
			(2)
	(ii)	Name Symbol	(2)
(b)	seei	dioactive source was placed within 2 cm of the spark counter and lots of sparks were n. A piece of paper was slid between the source and the counter. The sparking sped.	
	(i)	What type of radiation was being given off?	
			(4)
	(ii)	The paper was removed and the source slowly moved away from the spark counter. Describe what will happen to the sparking.	(1)
			(2)


(c)	A radioactive source gave a high reading using a Geiger-Müller tube and counter not cause sparking when brought near to the spark counter. Why?	r, but did
		(1) (Total 8 marks)

Some students measure the level of radiation from a radioactive source during the same lesson each week over a period of six weeks.

Here are the results. (They have been corrected for background radiation.)

Time (weeks)	start	1	2	3	4	5	6
Level of radiation (average counts per minute)	66	44	34	29	16	12	8

(a) Using the graph paper below, display these results in the most appropriate way.




(5)

www.tutorzone.co.t	What overall pattern is there in the students' results?	(b)
(3)		
(U) (Total & marks)		

Some rocks contain the radioactive isotope uranium–235 (235U).

<sup>235</sup>U has a half-life of 700 million years and, as it decays, lead–207 (<sup>207</sup>Pb) is eventually formed.

(a) Draw a decay curve for <sup>235</sup>U on the graph below.



(4)

Samples of an igneous rock gave an average ratio of 70 atoms of <sup>235</sup>U to 30 atoms (b) of 207 Pb.

Use the decay curve you have drawn to estimate the age of the igneous rock.

Answer ..... million years.

(1)

(c) A sandstone rock which lies above the igneous rock contains traces of uranium-235 and of lead-207.

Why might it be unsatisfactory to use this uranium for dating the sandstone?

.....

(Total 7 marks)

(a) When an atom of thorium-232 decays, an alpha (α) particle is emitted from the nucleus. An atom of radium is left behind.

An alpha particle consists of two protons and two neutrons.

We can represent this radioactive decay in a special kind of equation:

Th 
$$\xrightarrow{4}$$
 a  $\xrightarrow{228}$   $\xrightarrow{4}$  - - number of nucleons (protons + neutrons)

Ra

88  $\xrightarrow{4}$  - - number of protons

Thorium-228 is also radioactive.

Atoms of this isotope also decay by emitting an alpha particle and producing an isotope of radium.

Complete the equation for this decay.

58

(4)

(b) An atom of radium-228 decays by emitting a beta (β) particle from the nucleus.

A beta particle is in fact an electron (symbol  $_{-1}^{0}e$ ).

The effect of this is to change a neutron into a proton.

An atom of actinium remains.

This type of decay can also be represented by an equation:

Ra 
$$\longrightarrow$$
 e  $\stackrel{228}{-1}$  e  $\stackrel{228}{-1}$   $\stackrel{--}{-}$  number of nucleons (protons + neutrons)

This isotope of actinium is radioactive.

An atom of actinium-228 also decays by emitting a beta particle.


An isotope of thorium is left behind.

Complete the equation for this decay.

(4)

(c) Thorium-232 eventually decays to the stable isotope lead-208.

All the steps in this process can be shown on a diagram.



(i) Complete the sentences:

During the decay from (A) to (B) a ...... particle is emitted.

During the decay from (B) to (C) a ...... particle is emitted.

During the decay from (E) to (F) a ...... particle is emitted.

During the decay from (I) to (J) a ...... particle is emitted.

(2)

The table shows how long it takes for half of the atoms of each isotope to decay. (ii)

ISOTOPE	TIME FOR HALF TO DECAY	
A	billions of years	
В	7 years	
С	6 years	
D	2 years	
Е	4 days	
F	1 minute	
G	0.4 seconds	
Н	10 hours	
I	1 hour	
J	0.3 microseconds	

A rock sample contains:

- many atoms of thorium -232
- even more atoms of lead -208


<ul> <li>hardly any atoms of any of the other isotopes shown on the diagram</li> </ul>
Explain this as fully as you can.

(3) (Total 13 marks) Doctors sometimes need to know how much blood a patient has.

They can find out by using a radioactive solution.

After measuring how radioactive a small syringe-full of the solution is they inject it into the patient's blood.

## YOUR BLOOD CIRCULATION



They then wait for 30 minutes so that the solution has time to become completely mixed into the blood.

Finally, they take a syringe-full of blood and measure how radioactive it is.

## Example:

If the doctor injects  $10 \text{ cm}^3$  of the radioactive solution and this is diluted 500 times by the blood there must be  $10 \times 500 = 5000 \text{ cm}^3$  of blood.

(4)

After allowing for background radiation:	www.tutorzon
• 10 cm <sup>3</sup> of the radioactive solution gives a reading of 7350 counts per minutes.	ute;
• a 10 cm <sup>3</sup> sample of blood gives a reading of 15 counts per minute.	
Calculate the volume of the patient's blood. (Show your working.)	
	<ul> <li>10 cm<sup>3</sup> of the radioactive solution gives a reading of 7350 counts per minute.</li> <li>a 10 cm<sup>3</sup> sample of blood gives a reading of 15 counts per minute.</li> <li>Calculate the volume of the patient's blood.</li> </ul>




As time passes, the solution becomes less radioactive on its own. RADIATION 0 TIME (h)

Radiation from radioactive substances can harm your body cells.

(b)	The doctor's method of estimating blood volume will not be completely accurate. Write down <b>three</b> reasons for this.	
	1	
	2	
	3	(3)
(c)	The doctors use a radioactive substance which loses half of its radioactivity every six hours. Explain why this is a suitable radioactive substance to use.	

(Total 9 marks)

The diagram shows what happens to the radiation from three radioactive substances when different materials are put in the way.



Choose types of radiation from this list to complete the table below.

α (alpha)	β (beta)	γ ( <b>gamma</b> )	UV (ultraviolet)
RADIOACTIV	RADIOACTIVE SUBSTANCE		RADIATION IT EMITS
	A		
	В		
	С		

(Total 3 marks)